

Concepts: Algebraic combinations of functions, composition and decomposition of functions.

Algebraic Combinations of Functions

An ambitious way of creating new functions is to combine two or more functions to create a new function.

The most obvious way we can do this is to perform basic algebraic operations on the two functions to create the new one; hence we can add, subtract, multiply or divide functions.

Note that there are two types of algebras in use in this section,

1. the algebra of real numbers, i.e. $4 \times 5 = 20$, $4 - 5 = -1$, $20/10 = 2$, etc.,
2. the algebra of functions, fg , $f - g$, etc.

Algebra of functions

Let f (with domain A) and g (with domain B) be functions. Then the functions $f + g$, $f - g$, fg , f/g are defined as:

$$(f + g)(x) = f(x) + g(x) \text{ domain } A \cap B$$

$$(f - g)(x) = f(x) - g(x) \text{ domain } A \cap B$$

$$(fg)(x) = f(x)g(x) \text{ domain } A \cap B$$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} \text{ domain } \left\{x \in A \cap B \mid g(x) \neq 0\right\}$$

The domains are all the *intersection* (that's what the symbol \cap means) of the domain of f and g , making sure we don't divide by zero.

A Closer Look

- The minus sign in $f - g$ represents *the difference between two functions*.
- The minus sign in $f(x) - g(x)$ represents *the difference between two real numbers*.

The relation that we have that allows us to calculate this quantity is $(f - g)(x) = f(x) - g(x)$, which is easy to remember.

This is a subtle point, but it is always a good idea to understand what the mathematical notation is telling you.

Note: Two functions are equal if they have the same functional definition and the same domain.

Example If $f(x) = \sqrt{x}$ and $g(x) = \sqrt{4 - x^2}$, find the functions $f + g$, $f - g$, fg , f/g and give their domains.

First, we need to determine the intersection of the domains of f and g , so we need to determine the domains of f and g .

Domain of $f = \sqrt{x}$ is $x \in [0, \infty)$

Domain of $g = \sqrt{4 - x^2}$ is such that $4 - x^2 \geq 0 \rightarrow -2 \leq x \leq 2$ or $x \in [-2, 2]$.

Therefore the intersection of these domains is $x \in [0, 2]$ or $0 \leq x \leq 2$.

And our new functions are defined as:

$$(f + g)(x) = \sqrt{x} + \sqrt{4 - x^2}, 0 \leq x \leq 2$$

$$(f - g)(x) = \sqrt{x} - \sqrt{4 - x^2}, 0 \leq x \leq 2$$

$$(fg)(x) = \sqrt{x}\sqrt{4 - x^2} = \sqrt{4x - x^3}, 0 \leq x \leq 2$$

$$(f/g)(x) = \frac{\sqrt{x}}{\sqrt{4 - x^2}}, 0 \leq x < 2, \text{ where we exclude } x = 2 \text{ since it would lead to division by zero.}$$

Example If $f(x) = \sqrt{x}$, find the function ff and give the domain.

First, we need the domain of f : Domain of $f = \sqrt{x}$ is $x \in [0, \infty)$

Our new function is defined as

$$(ff)(x) = \sqrt{x}\sqrt{x} = x, 0 \leq x \leq \infty.$$

So the basic function $h(x) = x, x \in \mathbb{R}$ is NOT the same function as $(ff)(x) = x, 0 \leq x < \infty$ since the domains are different.

Composition of functions


Given two functions f and g , the *composite function* $f \circ g$ (called the composition of f and g) is defined by

$$(f \circ g)(x) = f(g(x))$$

and the domain of $f \circ g$ consists of all x -values in the domain of g that map to $g(x)$ values in the domain of f .

It is important to note that $f \circ g \neq g \circ f$.

Arrow Diagram of Composition $f \circ g$



The important thing to notice is the domain of the composition $f \circ g$ is a subset of the domain of g . To get the domain of a composition, do not simplify the composition at all, and look for the x values for which that expression is defined.

Example Find $(f \circ g \circ h)(x)$ if $f(x) = \frac{x}{x+1}$, $g(x) = x^{10}$, $h(x) = x+3$.

$$\begin{aligned} (f \circ g \circ h)(x) &= f(g(h(x))) \\ &= f(g(x+3)) \\ &= f((x+3)^{10}) \\ &= \frac{(x+3)^{10}}{(x+3)^{10} + 1} \end{aligned}$$

Since the domain and range of h is $x \in \mathbb{R}$, and the domain and range of g is $x \in \mathbb{R}$, there are no restrictions on the domain from the set we are drawing from. The only restriction arises from division by zero, but since $(x+3)^{10} + 1 = 0$ has no real valued solutions, the domain of $f \circ g \circ h$ is $x \in \mathbb{R}$.

Composition example If $f(x) = \sqrt{x}$ and $g(x) = \sqrt{4 - x^2}$, find $(f \circ g)(x)$ and $(g \circ f)(x)$ and give the domains.

Solution

To get the domain, do not simplify at all and look for any areas of the expression that would cause problems.

$$\begin{aligned}(f \circ g)(x) &= f(g(x)) \\ &= f(\sqrt{4 - x^2}) \\ &= \sqrt{(\sqrt{4 - x^2})} \text{ (this line is used to get domain)} \\ &= (4 - x^2)^{1/4}\end{aligned}$$

For domain, the expression $\sqrt{(\sqrt{4 - x^2})}$ is only defined if $4 - x^2 \geq 0$, which means $-2 \leq x \leq 2$. Therefore, the domain of $f \circ g$ is $x \in [-2, 2]$.

$$\begin{aligned}(g \circ f)(x) &= g(f(x)) \\ &= g(\sqrt{x}) \\ &= \sqrt{4 - (\sqrt{x})^2} \text{ (this line is used to get domain)} \\ &= \sqrt{4 - x}\end{aligned}$$

For domain, the expression $\sqrt{4 - (\sqrt{x})^2}$ is only defined if $x \geq 0$ and $4 - (\sqrt{x})^2 \geq 0$. This is a compound inequality (two inequalities must be satisfied). The second inequality simplifies to $x \leq 4$, so both are satisfied when $x \in [0, 4]$. The domain of $g \circ f$ is $[0, 4]$.

Note this is different than if we had just looked at the domain of $h(x) = \sqrt{4 - x}$, which is $x \leq 4$.

The lesson: the domain of compositions cannot be found simply by looking at the final function relation.

Example of Decomposing using composition Given $F(x) = \cos^2(x + 9)$, determine functions f, g, h so you can write $F(x)$ as a composition $(f \circ g \circ h)(x)$.

Look at how you compute $F(x)$, and build the functions from that:

Add 9: $h(x) = x + 9$

Take cosine: $g(x) = \cos x$

Square: $f(x) = x^2$

You should check that this choice yields $(f \circ g \circ h)(x) = F(x)$.