

Questions

Example (2.2.1) Solve the differential equation $y' = x^2/y$.

Example (2.2.4) Solve the differential equation $y' = (3x^2 - 1)/(3 + 2y)$.

Example (2.2.24) Solve the initial value problem $y' = (2 - e^x)/(3 + 2y), y(0) = 0$ and determine where the solution attains its maximum value.

Solutions

Example (2.2.1) Solve the differential equation $y' = x^2/y$.

$$\begin{aligned}\frac{dy}{dx} &= \frac{x^2}{y} \\ y \, dy &= x^2 \, dx \\ \int y \, dy &= \int x^2 \, dx \\ \frac{y^2}{2} &= \frac{x^3}{3} + c\end{aligned}$$

This is an implicit solution for y .

Example (2.2.4) Solve the differential equation $y' = (3x^2 - 1)/(3 + 2y)$.

$$\begin{aligned}\frac{dy}{dx} &= \frac{3x^2 - 1}{3 + 2y} \\ (3 + 2y) \, dy &= (3x^2 - 1) \, dx \\ \int (3 + 2y) \, dy &= \int (3x^2 - 1) \, dx \\ 3y + y^2 &= x^3 - x + c\end{aligned}$$

This is an implicit solution for y .

Example (2.2.24) Solve the initial value problem $y' = (2 - e^x)/(3 + 2y), y(0) = 0$ and determine where the solution attains its maximum value.

$$\begin{aligned}\frac{dy}{dx} &= \frac{2 - e^x}{3 + 2y} \\ (3 + 2y) \, dy &= (2 - e^x) \, dx \\ \int (3 + 2y) \, dy &= \int (2 - e^x) \, dx \\ 3y + y^2 &= 2x - e^x + c\end{aligned}$$

We use the initial condition $y(0) = 0$ to determine the constant c . When $x = 0, y = 0$:

$$\begin{aligned}0 &= -1 + c \\ c &= 1\end{aligned}$$

The implicit solution to the initial value problem is $3y + y^2 = 2x - e^x + 1$.

The maximum occurs when $y' = 0$. Implicitly differentiate:

$$\begin{aligned}\frac{d}{dx}[3y + y^2] &= 2x - e^x + c \\ 3\frac{dy}{dx} + 2y\frac{dy}{dx} &= 2 - e^x \\ 0 &= 2 - e^x \\ e^x &= 2 \\ x &= \ln 2\end{aligned}$$

This is where the maximum occurs.
